

## The Negative Binomial Distribution

Study Notes | Written by Larry Cui

Follow the logic of geometric distribution, if we want to study the probability of  $r^{th}$  success in  $k^{th}$  of a series of trials, it must be the case that  $(r - 1)$  success occur during the first  $(k - 1)$  trials and the  $r^{th}$  happens on exactly the  $k^{th}$  trial.

### Negative Binomial Distribution pdf

$$p_X(k) = \binom{k-1}{r-1} p^r (1-p)^{k-r}, \quad k = r, r+1, \dots$$

If we let  $X$  be the sum of independent variables  $X_1, X_2, \dots, X_r$ , and let  $X \rightarrow \infty$ , the negative binomial can be interpreted as  $r$  successes happen one after another, and each of which follows the geometric distribution model. This perspective won't give us a better form to the pdf, but will greatly simplify the calculation of expected values.

### E(X), Var(X) and mgf

Let  $p_X(k) = \binom{k-1}{r-1} p^r (1-p)^{k-r}$ ,  $k = r, r+1, \dots$  then:

- $M_X(t) = M_{X_1}(t)M_{X_2}(t) \dots M_{X_r}(t) = \left[ \frac{pe^t}{1 - (1-p)e^t} \right]^r$
- $E(X) = E(X_1) + E(X_2) + \dots + E(X_r) = \frac{r}{p}$
- $\text{Var}(X) = \text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_r) = \frac{r(1-p)}{p^2}$

### Algebraic Approach: the sum of pdf

There's an algebraic approach to the  $E(X)$  and  $\text{Var}(X)$ , which centres around the verification of the pdf of the function. Let's take a look at the following lemma:

#### lemma (a)

$$\binom{k+m}{m} = \binom{k+m-1}{m} + \binom{k+m-1}{m-1}$$

**Proof:** a simple expansion of the combinations gives us result. Furthermore, we can also see this equation by some intuition: if you want pick  $m$  items from  $k+m$  pool, you can first

split the pool into one item and the other. The final pick consists of two scenarios: 1) excluding this one item:  $\binom{k+m-1}{m}$ ; and 2) including this one item:  $\binom{k+m-1}{m-1}$ .

$$\begin{aligned}\binom{k+m-1}{m} + \binom{k+m-1}{m-1} &= \frac{(k+m-1)!}{m!(k-1)!} + \frac{(k+m-1)!}{(m-1)!k!} \\ &= \frac{(k+m-1)!k + (k+m-1)!m}{m!k!} \\ &= \frac{(k+m)!}{m!k!}\end{aligned}$$

**Intermediate function:** consider the function

$$f_m(z) = \sum_{k=0}^{\infty} \binom{k+m}{m} z^k$$

Use lemma (a), we can expand it as

$$f_m(z) = \sum_{k=0}^{\infty} \binom{k+m-1}{m-1} z^k + \sum_{k=0}^{\infty} \binom{k+m-1}{m} z^k$$

Because  $\binom{m-1}{m} = \frac{(m-1)(m-2)\cdots(m-m)}{m!(-1)!} = 0$  by definition, we can discard the  $k=0$  item from the second term. A little re-arrangement give the following form for the second term:

$$\sum_{k=1}^{\infty} \binom{k+m-1}{m} z^k = z \sum_{k=1}^{\infty} \binom{k+m-1}{m} z^{k-1} = z f_m(z)$$

so

$$f_m(z) = f_{m-1}(z) + z f_m(z) \quad \Rightarrow \quad f_m(z) = \frac{f_{m-1}(z)}{1-z}$$

But since  $f_0(z) = \sum_{k=0}^{\infty} \binom{k}{0} z^k = \frac{1}{1-z}$ , we know  $f_m(z) = \frac{1}{(1-z)^{m+1}}$ .

To verify the sum of the pdf equals to 1, we use the above function to get (here we let  $m = r-1, k' = k-r$ ):

$$\begin{aligned}\sum_{k=r}^{\infty} \binom{k-1}{r-1} p^r (1-p)^{k-r} &= p^r \cdot \sum_{k'=0}^{\infty} \binom{k'+m}{m} (1-p)^{k'} \\ &= p^r \cdot f_m(1-p) \\ &= p^r \cdot \frac{1}{(1-1+p)^r} = 1\end{aligned}\tag{1}$$

Remark:  $r = m+1$

Now we can go on with  $E(X)$ :

$$\begin{aligned}
 E(X) &= \sum_{k=r}^{\infty} \binom{k-1}{r-1} k p^r (1-p)^{k-r} \\
 &= p^r \cdot \sum_{k'=0}^{\infty} (k'+r) \binom{k'+m}{m} (1-p)^{k'} \\
 &= p^r \cdot \sum_{k'=0}^{\infty} r \binom{k'+r}{r} (1-p)^{k'} \quad \text{Remark: } m = r-1 \\
 &= r p^r \cdot \frac{1}{(1-1+p)^{r+1}} = \frac{r}{p}
 \end{aligned} \tag{2}$$

In order to find  $\text{Var}(X)$ , we need a little trick here: find  $E[X(X+1)]$  first. As above, we start by lay out the formula:

$$\begin{aligned}
 E[X(X+1)] &= \sum_{k=r}^{\infty} k(k+1) \binom{k-1}{r-1} p^r (1-p)^{k-r} \\
 &= p^r \cdot \sum_{k'=0}^{\infty} (k'+r+1)(k'+r) \binom{k'+m}{m} (1-p)^{k'} \quad \text{Remark: } k = k' + r \\
 &= p^r \cdot \sum_{k'=0}^{\infty} (k'+r+1)(k'+r) \binom{k'+r-1}{r-1} (1-p)^{k'} \quad \text{Remark: } m = r-1 \\
 &= r(r+1)p^r \cdot \sum_{k'=0}^{\infty} \binom{k'+r+1}{r+1} (1-p)^{k'} \\
 &= r(r+1)p^r \cdot \frac{1}{(1-1+p)^{r+2}} \\
 &= \frac{r(r+1)}{p^2}
 \end{aligned}$$

so we have

$$\text{Var}(X) = E[X(X+1)] - E(X)^2 - E(X) = \frac{r(r+1)}{p^2} - \frac{r^2}{p^2} - \frac{r}{p} = \frac{r(1-p)}{p^2}$$

### A second way:

We can also find the solution to Equation (1) without having recourse to intermediate function  $f_m(z)$ . First of all, we re-arrange the expected value formula as:

$$\begin{aligned}
 E(X) &= \sum_{k=r}^{\infty} \binom{k-1}{r-1} k p^r (1-p)^{k-r} \\
 &= p^r \cdot \sum_{k'=0}^{\infty} (k'+r) \binom{k'+r-1}{r-1} (1-p)^{k'} \quad \text{Remark: } k' = k - r \\
 &= r p^r \cdot \sum_{k'=0}^{\infty} \binom{k'+(r+1)-1}{k'} (1-p)^{k'} \\
 &= r p^r \cdot \sum_{k'=0}^{\infty} \binom{k'+(r+1)-1}{k'} (-1)^{k'} (p-1)^{k'}
 \end{aligned}$$

We need Taylor/Maclaurin series for help here. Recall

**The Maclaurin series for  $f(x)$**

wherever it converges, can be expressed as:

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \cdots + \frac{f^{(k)}(0)}{k!}x^k + \cdots$$

and for  $f(x) = 1/(1+x)^n$ :

$$\frac{1}{(1+x)^n} = 1 - nx + \frac{(-n)(-n-1)}{2!}x^2 + \cdots = \sum_{k=0}^{\infty} \binom{n+k-1}{k} (-1)^k x^k$$

We can see that the last part of  $E(X)$  is exactly the sum of Maclaurin series with  $n = r + 1$  and  $x = p - 1$ , so

$$E(X) = rp^r \cdot \frac{1}{(1+p-1)^{r+1}} = \frac{r}{p}$$